

Unravelling abiogenic and biogenic sources of methane in the Earth's deep subsurface

B. Sherwood Lollar ^{a,*}, G. Lacrampe-Couloume ^{a,1}, G.F. Slater ^{a,1}, J. Ward ^{a,1},
D.P. Moser ^{b,2}, T.M. Gihring ^{b,3}, L.-H. Lin ^{c,4}, T.C. Onstott ^{c,4}

^a Department of Geology, 22 Russell St., University of Toronto, Toronto, Ontario, Canada M5S 3B1

^b Environmental Microbiology Group, Pacific Northwest National Laboratory, Richland WA 99352, USA

^c Dept. of Geosciences, Guyot Hall, Princeton University, Princeton NJ 08544, USA

Accepted 5 September 2005

Abstract

At four underground sites in Precambrian Shield rocks in Canada and South Africa, hydrocarbon and hydrogen gases exsolving from saline fracture waters are analyzed for compositional and isotopic signatures. Dominated by reduced gases such as CH₄, H₂ and higher hydrocarbons (ethane, propane, butane), the most ¹³C-enriched methane end-members at all four sites show a pattern of carbon and hydrogen isotopic values similar to abiogenic gases produced by water–rock interaction that have been identified previously at one site on the Precambrian Shield in Canada. The abiogenic nature of these gases was not previously recognized due to mixing with a second methane component produced by microbial processes. The microbial methane end-member is identified based on carbon and hydrogen isotopic signatures, and DNA gene amplification (PCR) data that indicate the presence of methanogens. A framework is presented to estimate the relative contribution of abiogenic versus microbial hydrocarbon gases at these sites. This approach has important implications for evaluation of potential abiogenic hydrocarbon reservoirs in a wide range of geologic settings, including the longstanding controversy concerning the possible contribution of abiogenic gases to economic petroleum hydrocarbon reservoirs. The association of high concentrations of H₂ with ¹³C-enriched CH₄ end-members, and H₂ depletion in the ¹³C-depleted methanogenic end-members further suggests the possibility that abiogenic gases may support H₂ autotrophy linked to methanogenesis in the deep subsurface.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Abiogenic; Methane; Hydrogen; Autotrophy; Deep biosphere; Mars

* Corresponding author. Tel.: +1 416 978 0770; fax: +1 416 978 3938.

E-mail addresses: bsllollar@chem.utoronto.ca (B. Sherwood Lollar), glc@geology.utoronto.ca (G. Lacrampe-Couloume), gslater@mcmaster.ca (G.F. Slater), ward@geology.utoronto.ca (J. Ward), duane.moser@dri.edu (D.P. Moser), gihring@ocean.fsu.edu (T.M. Gihring), lhlin@ntu.edu.tw (L.-H. Lin), tullis@princeton.edu (T.C. Onstott).

¹ Fax: +1 416 978 3938.

² Fax: +1 702 862 5360.

³ Fax: +1 850 644 2581.

⁴ Fax: +1 609 258 1274.

1. Introduction

Recent reports of CH₄ in the Mars atmosphere focussed scientific and public attention on possible geological and biological sources of these hydrocarbons (Formisano et al., 2004; Kerr, 2004). Resolving the question of the origin of atmospheric CH₄ on Mars is even more challenging given that distinguishing abiogenic versus biogenic sources of CH₄ in the terrestrial subsurface is still controversial (Gold, 1979; Kenney et al., 2002; Shock, 1995). Deep subsurface fluids in Precambrian Shield rocks have been shown to be dominated by reduced gases such as CH₄ and locally, high concentrations of H₂ (up to 30% by volume) (Sherwood Lollar et al., 1993a,b). Sherwood Lollar et al. (2002) used stable isotope signatures to suggest that CH₄ and higher hydrocarbon gases (ethane, propane and butane) at Kidd Creek mine on the Canadian Shield are produced abiogenically by water–rock interaction such as surface-catalysed polymerization (Anderson, 1984; Foustoukos and Seyfried, 2004); metamorphism of graphite–carbonate bearing rocks (Giardini and Salotti, 1969; Holloway, 1984; Kenney et al., 2002); and other gas–water–rock alteration reactions such as serpentinization (Berndt et al., 1996; Charlou and Donval, 1993; Horita and Berndt, 1999; Kelley et al., 2001, 2005; McCollom and Seewald, 2001; Vanko and Stakes, 1991). In this paper, data from 4 new sites in Canada and South Africa suggests that abiogenic hydrocarbon gases are more globally pervasive than has been understood previously. This is due to the fact that at many sites, the distinct abiogenic signature of such hydrocarbons is obscured by mixing with microbial CH₄. Based on this data, we present a model for identification of abiogenic hydrocarbons through the resolution of microbial and abiogenic mixing — an approach that is applicable to a wide variety of crustal settings where potential abiogenic hydrocarbon reserves have been suggested (Gold, 1979; Kenney et al., 2002; Shock, 1995).

2. Geological setting and samples

In gold and base-metal mines throughout the Precambrian Shield rocks of Canada, Finland and South Africa, flammable gases discharge from fractures and exploration boreholes (Nurmi and Kukkonen, 1986; Lahermo and Lampen, 1987; Cook, 1998; Sherwood Lollar et al., 1993a,b). Originally dissolved in saline groundwater (TDS levels range from several 1000 to tens of 1000s of ppm) in sealed fracture systems in the rocks, gases are released via depressurization into mine

workings at rates of 1 to >30 L gas/min/borehole (Sherwood Lollar et al., 1993a,b). The groundwaters are typically NaCaCl-rich or CaNaCl-rich with low levels of SO₄ (typically <15 ppm and always <100 ppm) with pH levels between 7–9. This study focuses on two sites in Canada (Kidd Creek and Copper Cliff South mines) and three in South Africa (Driefontein, Kloof and Mponeng mines). Kidd Creek mine, situated in the southern volcanic zone of the Abitibi greenstone belt (approximately 2700 Ma), 24 km N of Timmins, Ontario is one of the world's largest volcanogenic massive sulfide deposits. All samples were collected from 2–2.1 kmbls (kilometers below land surface). The Cu–Ni ore at Copper Cliff South mine (CCS) in Sudbury, Ontario is associated with a quartz diorite dike offset of the Sudbury Igneous Complex — a micropegmatite, norite and quartz diorite irruptive emplaced approximately 1840 Ma (Cochrane, 1984) and samples were from approximately 1.3 kmbls. In South Africa, Kloof, Driefontein and Mponeng mines are all located west of Johannesburg in the Witwatersrand Basin — a large Archean intracratonic basin composed of volcanosedimentary sequences that unconformably overlie 3.0 Ga granitic basement. The basin is divided chronologically into quartzite, shale and minor volcanic strata of the Witwatersrand Supergroup (2900 Ma), the basaltic lava sequence of the Ventersdorp Supergroup (2700 Ma) and the overlying clastic and dolomitic sediments of the Transvaal Supergroup (2400–2500 Ma) (Coward et al., 1995). Almost all the samples in this study are from the Ventersdorp Supergroup and are located between 2.7 and 3.4 kmbls. The exceptions are borehole DR938H3, which while drilled in the Ventersdorp, intersects the underlying Witwatersrand Supergroup at approximately half its 750 m length, and DR9IPC, which is drilled in the overlying Transvaal Supergroup and is located at 0.9 kmbls.

3. Methodology

3.1. Sampling methods

All gas and fracture water samples were collected at the borehole collar after the method of Sherwood Lollar et al. (2002) and Ward et al. (2004). A packer was placed into the opening of the borehole and sealed to the inner rock walls below water level to seal the borehole from the mine air and minimize air contamination. Gas and water were allowed to flow through the apparatus long enough to displace any air remaining in the borehole or the apparatus before sampling. Plastic tubing was attached to the end of the packer and the

flow of gas and/or water from the borehole was directed into an inverted graduated funnel. Gases collected in the inverted funnel were transferred directly into evacuated vials through a needle that was attached to the top of the funnel. The gas sampling vials were pre-evacuated 130 ml borosilicate vials sealed with butyl blue rubber stoppers prepared after the method of Oremland and Des Marais (1983). Vials were pre-fixed with 50 μ l of a saturated HgCl₂ solution to kill any microbes contained in the sample so microbial activity post-sampling would not alter the gas composition and isotopic signatures. Previous studies, comparing the isotopic values of gases taken at the borehole collars to values determined for gases in solution at depth in the same boreholes, showed that exsolution of the hydrocarbon gases from solution does not alter their isotopic signatures (Sherwood Lollar et al., 1993a,b, 1994).

For the South African sites, the packer was sterilized by autoclaving before insertion into the borehole in order to reduce the possibility of introducing surface microorganisms by that means. In mining environments, aseptic drilling techniques were not possible. Microbial populations were characterized and compared however between the saline fracture waters from these sites, the low salinity mine service water, and ventilation air. Comparison of 16S rRNA gene clone libraries from these possible sources of contamination and from the fracture waters yielded no commonalities (Kieft et al., 1999; Onstott et al., 2003; Takai et al., 2001). In addition the fact that dissimilatory sulphate reductase (DSR) and 16S ribosomal (rRNA) genes amplified from the fracture water DNA yield sequences that are most similar to those reported for sulphate-reducing bacteria isolated from other subsurface sites is consistent with the fracture water microbial populations being in situ subsurface communities and not surface contaminants (Baker et al., 2003).

3.2. Compositional gas analysis

Compositional analyses of gas samples were performed at the Stable Isotope Laboratory at the University of Toronto. A Varian 3400 GC equipped with a flame ionization detector (FID) was used to determine concentrations of CH₄, C₂H₆, C₃H₈ and C₄H₁₀. The hydrocarbons were separated on a J and W Scientific GS-Q column (30 m \times 0.32 mm ID) with a helium gas flow and temperature program: initial 60 °C hold 2.5 min, increase to 120 °C at 5 °C/min. A Varian 3800 GC equipped with a micro-thermal conductivity detector (μ TCD) and a Varian Molecular Sieve 5A PLOT column (25 m \times 0.53 mm ID) was used to determine

concentrations of the inorganic gas components (H₂, He, Ar, O₂, CO₂ and N₂). To determine concentrations of Ar, O₂ and N₂ the helium gas flow rate was 3 ml/min and the temperature program was: initial 30 °C hold 6 min, increase to 80 °C at 15 °C/min, hold 4 min. To determine CO₂ concentrations, the helium gas flow rate was 50 ml/min and temperature program was: initial 60 °C, increase to 250 °C at 20 °C/min, hold 6 min. To determine concentrations of H₂ and He, the argon carrier gas flow rate was 2 ml/min and temperature program was: initial 10 °C hold 10 min, increase to 80 °C at 25 °C/min, hold 7 min. All analyses were run in triplicate and mean values are reported in Table 1. Reproducibility for triplicate analyses was \pm 5%.

3.3. Isotopic analysis

Stable carbon and hydrogen isotopic analysis for all hydrocarbons were performed at the University of Toronto (Table 2). Analyses for $\delta^{13}\text{C}$ values were performed by continuous flow compound specific carbon isotope ratio mass spectrometry with a Finnigan MAT 252 mass spectrometer interfaced with a Varian 3400 capillary GC. Hydrocarbons were separated by a Poraplot QTM column (25 m \times 0.32 mm ID) with temperature program: initial 40 °C hold 1 min, increase to 190 °C at 5 °C/min, hold 5 min. To separate CO₂ from CH₄ the program was started at 10 °C (hold 2 min) increased to 190 °C at 5 °C/min (hold 5 min). Total error incorporating both accuracy and reproducibility is \pm 0.5‰ with respect to V-PDB standard.

The $\delta^2\text{H}$ analysis was performed on a continuous flow compound specific hydrogen isotope mass spectrometer which consists of an HP 6890 gas chromatograph (GC) interfaced with a micropyrolysis furnace (1465 °C) in line with a Finnigan MAT Delta⁺-XL isotope ratio mass spectrometer. The hydrocarbon gases were separated on a Poraplot QTM column (25 m \times 0.32 mm ID) with a helium carrier at 2.2 ml/min and temperature program: initial 35 °C hold 3 min, increase to 180 °C at 15 °C/min. Total error incorporating both accuracy and reproducibility is \pm 5‰ with respect to V-SMOW.

3.4. DNA gene amplification (PCR)

Methyl coenzyme M reductase α -subunit genes (*mcrA*) were amplified from community DNA extracts according to the methods of Hales et al. (1996). Archaeal 16S rRNA gene was amplified using a nested PCR reaction in which PCR product generated using the primers 21F (Moyer et al., 1998), and 1492R (Rey-

Table 1
Gas compositional data (in %)

Site	Borehole No.	Ar	H ₂	He	O ₂	N ₂	CH ₄	C ₂ H ₆	C ₃ H ₈	C ₄ H ₁₀
Copper Cliff	CCS4577	0.06	54.0	3.46	0.11	1.92	34.4	5.31	0.49	0.11
Copper Cliff	CCS4547	0.07	43.0	3.38	0.12	2.20	38.1	5.28	0.56	0.12
Copper Cliff	CCS4546	0.17	9.94	4.37	0.52	6.38	69.5	7.14	0.78	0.17
Copper Cliff	CCS4572	0.05	57.8	2.62	0.15	1.72	25.3	4.16	0.38	0.09
Copper Cliff	CCS4880	0.27	19.7	6.42	0.10	4.54	59.3	7.39	1.11	0.29
Mponeng	MPA	NA	3.30	9.06	4.77	28.1	53.8	2.47	0.36	0.08
Mponeng	MP104	NA	11.5	12.3	3.65	21.1	49.6	3.99	0.60	0.13
Driefontein	DR548	NA	10.3	3.05	6.18	23.6	50.7	3.85	0.52	0.11
Driefontein	DR938H1	NA	<0.01	0.90	16.5	72.5	5.45	0.14	0.02	<0.01
Driefontein	DR938H3-0m	NA	0.74	5.98	0.55	14.0	76.0	3.15	0.32	0.06
Driefontein	DR938H3'-0m*	NA	0.32	4.64	4.95	28.6	61.4	2.45	0.26	0.04
Driefontein	DR938H3-648 m	NA	NA	NA	NA	NA	NA	NA	NA	NA
Driefontein	DR938CH1	NA	<0.05	0.21	23.4	89.9	1.19	<0.05	<0.05	<0.05
Driefontein	DR9IPC	NA	<0.05	2.75	3.77	81.9	11.6	<0.05	<0.05	<0.05
Kloof	KL1GH	NA	9.15	3.42	14.5	56.4	10.7	0.49	<0.01	<0.01
Kloof	KL441H3	NA	<0.01	15.9	0.43	45.1	33.0	0.66	0.07	0.02
Kloof	KL739	NA	9.25	13.5	<0.04	7.72	64.9	2.86	0.41	0.08
Kloof	KL441H2	1.68	<0.01	15.6	6.55	61.9	23.4	0.51	0.05	<0.01
Kloof	KL443HWND1	1.39	<0.01	18.2	1.24	13.3	57.3	2.03	0.28	0.05
Kloof	KL443HWND	1.79	<0.01	20.8	0.97	16.8	53.4	2.04	0.21	0.03

a. All CO₂ was below detection limit.

b. NA — not analyzed.

c. C₄H₁₀ incorporates both *n*-butane and *iso*-butane.

d. *DR938H3'-0 m is a sample from the same borehole as DR938H3-0 m taken 8 months afterwards. Such temporal sampling was not possible for the other boreholes due to the common practice of sealing them immediately after completion of drilling.

e. DR938H3-0 m and DR938H3-648 m were collected from the same borehole on the same day. The 0 m sample was taken at the borehole collar while the 648 m sample was taken at 648 m depth.

senbach et al., 2000) was used as template in PCR reactions with the “nested” primers 21F and 958R (Moyer et al., 1998). *McrA* and archaeal 16S rRNA gene PCR products were cloned using Invitrogen TOPO kits. For each library, 48 clones were screened by restriction fragment length polymorphism. Unique clones were sequenced, then analyzed using BLAST (Altschul et al., 1997) to identify sequences in the NCBI Genbank database having high similarity (Table 3). Sequence identities were calculated using BioEdit.

4. Results and discussion

4.1. Isotopic patterns suggest an abiogenic origin

Mantle-derived abiogenic hydrocarbons are typically identified based on three criteria: a $\delta^{13}\text{C}$ value for CH₄ more enriched than -25% ; a “carbon isotopic reversal” trend of increasing isotopic depletion in ^{13}C with increasing molecular weight for CH₄–ethane–propane–butane; and a $^3\text{He}/^4\text{He}$ ratio indicative of mantle-derived helium ($R/R_a > 0.1$) (Jenden et al., 1993). Based on these criteria, coal bed or thermogenic hydrocarbon gases with anomalously enriched $\delta^{13}\text{C}$ values and/or anomalously depleted $\delta^2\text{H}$ values for methane could be

distinguished from mantle-derived abiogenic gases (Jenden et al., 1988; Jenden and Kaplan, 1989; Poreda et al., 1986, 1988; Lyon and Hulston, 1984; Rigby and Smith, 1981). Over the past decade however, there has been a growing body of literature that indicates that not all abiogenic gases are mantle-derived. A variety of low temperature water–rock interactions have been shown to produce both CH₄ and higher hydrocarbons such as ethane, propane, and butane — including surface-catalyzed polymerization from reduction of CO or CO₂ in a Fischer–Tropsch synthesis; heating or metamorphism of graphite or carbonate bearing rocks; or other vapor–water–rock alteration reactions such as serpentinization (Holloway, 1984; Yuen et al., 1990; Berndt et al., 1996; Hu et al., 1998; Horita and Berndt, 1999; McCollom and Seewald, 2001; Foustoukos and Seyfried, 2004). Significantly, several of these studies have demonstrated that production of abiogenic CH₄ by these water–rock interactions can result in $\delta^{13}\text{C}$ values as depleted as -57% (Yuen et al., 1990; Hu et al., 1998; Horita and Berndt, 1999). In a crustal-dominated geologic setting, these processes will produce abiogenic hydrocarbons whose $\delta^{13}\text{C}$ values reflect local crustal carbon sources and will not have either ^{13}C -enriched $\delta^{13}\text{C}$ values or R/R_a values > 0.1 . In addition, it has

Table 2
Carbon and hydrogen isotopic values (in ‰) for CH₄–C₄ hydrocarbon gases

Site	Borehole no.	$\delta^{13}\text{C}_{\text{CH}_4}$	$\delta^2\text{H}_{\text{CH}_4}$	$\delta^{13}\text{C}_{\text{C}_2}$	$\delta^2\text{H}_{\text{C}_2}$	$\delta^{13}\text{C}_{\text{C}_3}$	$\delta^2\text{H}_{\text{C}_3}$	$\delta^{13}\text{C}_{\text{C}_n\text{C}_4}$	$\delta^2\text{H}_{\text{C}_n\text{C}_4}$
Copper Cliff	CCS4577	–36.1	–413	–24.6	–266	ND	ND	ND	ND
Copper Cliff	CCS4547	–34.4	–424	–28.1	–290	ND	ND	ND	ND
Copper Cliff	CCS4546	–32.0	–452	–34.6	–348	–33.4	ND	ND	ND
Copper Cliff	CCS4572	–36.7	–401	–24.4	–269	ND	ND	ND	ND
Copper Cliff	CCS4880	–37.3	–372	–31.6	–301	ND	ND	ND	ND
Mponeng	MPA	–27.8	–349	–35.0	–267	–33.3	–212	ND	ND
Mponeng	MP104	–32.8	–366	–37.6	–270	–34.8	–193	–35.3	–88
Driefontein	DR548	–46.5	–403	–50.5	–285	–47.2	–192	–45.6	–72
Driefontein	DR938H1	–53.6	–295	–42.9	–326	–42.3	ND	ND	ND
Driefontein	DR938H3-0 m	–40.2	–368	–43.1	–290	–42.0	ND	ND	ND
Driefontein	DR938H3'-0 m*	–42.3	–369	ND	ND	ND	ND	ND	ND
Driefontein	DR938H3-648 m	–40.9	–371	–44.0	–234	–42.9	–179	ND	ND
Driefontein	DR938CH1	–47.5	–329	–43.2	ND	ND	ND	ND	ND
Driefontein	DR9IPC	–55.5	–218	ND	ND	ND	ND	ND	ND
Kloof	KL1GH	–29.2	–312	–33.2	–230	ND	ND	ND	ND
Kloof	KL441H3	–40.0	–253	–30.9	–241	–25.2	ND	ND	ND
Kloof	KL739	–28.7	–300	–30.7	–230	–27.2	–142	ND	ND
Kloof	KL441H2	–37.8	–257	–32.9	–198	–34.0	–105	ND	ND
Kloof	KL443HWDN1	–34.4	–327	–35.6	–251	–33.7	–208	ND	ND
Kloof	KL443HWDN	–34.3	–329	–34.3	–269	–33.5	–206	–33.6	–207

a. ND — below detection limit for isotope analysis.

b. *DR938H3'-0m is a sample from the same borehole as DR938H3-0m taken 8 months afterwards. Such temporal sampling was not possible for the other boreholes due to the common practice of sealing them immediately after completion of drilling.

c. DR938H3-0m and DR938H3-648m were collected from the same borehole on the same day. The 0 m sample was taken at the borehole collar while the 648 m sample was taken at 648 m depth.

become clear that the “carbon isotopic reversal” trend alone is not sufficient evidence to support an abiogenic origin (Horita and Berndt, 1999).

Based on crustal ³He/⁴He ratios at all sites, no mantle-derived component for the Precambrian Shield hydrocarbon gases is indicated (Lippmann et al., 2003; Sherwood Lollar et al., 1993b). Sherwood Lollar et al. (2002) suggested a crustal abiogenic origin due to water–rock interactions for the hydrocarbon gases at Kidd Creek however, based on a new criteria — an inverse correlation of ¹³C-depletion and ²H-enrichment between CH₄ and ethane (Fig. 1A). Building on a hypothesis for carbon isotope distribution patterns first proposed by Des Marais et al. (1981), it was suggested that this pattern results from synthesis of higher molecular weight hydrocarbons from CH₄ during polymerization. In such a process ¹²CH₄ reacts faster than ¹³CH₄ to form chains, so that ¹²C is more likely to be incorporated

into larger hydrocarbon chains in polymerization reactions, whereas, owing to preferential cleavage of the weaker ¹²C–¹H bond versus the ¹²C–²H bond, the light (¹H) isotope will be preferentially eliminated (Sherwood Lollar et al., 2002). In contrast, the pattern for commercial thermogenic natural gas deposits consists of a positive correlation of $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values between CH₄, ethane and the higher hydrocarbon gases propane and butane (Fig. 1A). This isotopic pattern of increasing isotopic enrichment in both ¹³C and ²H with increasing molecular weight for the C₁–C₄ homologues for thermogenic gases results from production of hydrocarbons by thermal cracking of a high molecular weight organic precursor (Des Marais et al., 1981; Schoell, 1988). At each of the 4 newly investigated Precambrian Shield sites in Canada and South Africa from this study, hydrocarbon gases were identified with the same pattern of ¹³C-depletion and ²H-enrichment between CH₄ and

Table 3
16S rRNA gene and *mcrA* clones from borehole DR9IPC having high relatedness to known methanogenic organisms

DR9IPC clone (Genbank accession number)	Percent of library	Nearest cultivated relative	Identity (%)
16S rRNA 1 (AY604061)	40.0	<i>Methanobacterium curvum</i>	93
<i>mcrA</i> 1 (AY604057)	23.7	<i>Methanosarcina barkeri</i>	99
<i>mcrA</i> 2 (AY604058)	73.7	<i>Methanobacterium bryantii</i>	86
<i>mcrA</i> 3 (AY604059)	2.7	<i>Methanobacterium aarhusense</i>	83




Fig. 1. Plot of $\delta^{13}\text{C}$ versus $\delta^2\text{H}$ values for C_1 – C_4 for Kidd Creek (A) and for 4 new Precambrian Shield sites (B). In the lower panel, the most ^{13}C -enriched samples at each site — Driefontein (open squares); Mponeng (closed circles), Kloof (closed triangles) and CCS (crosses) all show the characteristic ^{13}C depletion and ^2H enrichment between CH_4 and ethane similar to the proposed abiogenic gases at Kidd Creek (data from (Sherwood Lollar et al., 2002) – closed squares– upper panel). In marked contrast, the pattern for thermogenic gas reservoirs is a positive correlation of $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values for CH_4 and ethane due to increasing isotopic enrichment in both ^{13}C and ^2H with increasing molecular weight for the C_1 – C_4 homologues (open circles — upper panel). This pattern results from production of the hydrocarbon gases by thermal cracking of a high molecular weight organic precursor (Des Marais et al., 1981; Schoell, 1988). Error bars on $\delta^{13}\text{C}$ values are $\pm 0.5\text{‰}$. Error bars on both A and B on $\delta^2\text{H}$ values are $\pm 5\text{‰}$ and are smaller than the plotted symbols.

ethane as the proposed abiogenic gases from Kidd Creek (Fig. 1B; Table 2). Interestingly, for each site only the most ^{13}C -enriched samples show this pattern.

4.2. Mixing trends

Whereas the Kidd Creek data set is remarkably uniform, one sample (KC7792) has a $\delta^{13}\text{C}_{\text{CH}_4}$ value significantly more depleted in ^{13}C and enriched in ^2H than other samples from this site (Fig. 1A). Sherwood Lollar et al. (2002) suggested that this might be due to mixing with a more ^{13}C -depleted and ^2H -enriched microbial CH_4 component in this borehole. In the current study, the hydrocarbon gas samples analysed for 4 new Precambrian Shield sites show a much wider range of CH_4 isotopic signatures than at Kidd Creek (Figs. 1B

and 2). In all cases, the proposed abiogenic signature was observed only for the samples with the most ^{13}C -enriched CH_4 values at each site (DR938H3 at both 0 and 685 m, DR548; KL739 and KL1GH; CCS4546; and both samples from Mponeng — MP104 and MPA; Fig. 1B). The distinct ^{13}C – ^2H isotopic patterns between CH_4 and ethane for these samples (Fig. 1B), as well as their significantly depleted $\delta^2\text{H}$ values compared to typical thermogenic gases (Fig. 2), do not support a thermogenic origin for these ^{13}C -enriched end-members. Furthermore, as at Kidd Creek, the majority of the data points from CCS, Driefontein and Kloof Mines fall along linear trends between the most ^{13}C -enriched and ^2H -depleted end-members, and samples with more ^{13}C -depleted and ^2H -enriched values (Fig. 2), raising the possibility of mixing between these two end-members. For CCS, Kloof and Kidd Creek, all data fall along a single trend from the ^{13}C -enriched end-members identified in Fig. 1B, whereas for Driefontein, two possible trends can be drawn through the data, one for DR938H3 and one for DR548. At Mponeng, both samples MP104 and MPA are ^{13}C -enriched and show the proposed abiogenic signature (Fig. 1B) and this is consistent with the lack of any mixing trend in Fig. 2.

In Fig. 2, solid lines indicate the conventionally accepted fields of $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values for microbial and thermogenic CH_4 based on empirical studies (Schoell, 1988). The hatched trend lines indicate that ^{13}C -depleted and ^2H -enriched end-members at these sites could theoretically be either thermogenic or microbial in origin based on the distribution of the $\delta^{13}\text{C}$ and $\delta^2\text{H}$ fields. Other lines of evidence indicate that this end-member is microbial however, and suggest that microbial CH_4 mixing with abiogenic gases produces the observed trend lines. For borehole DR9IPC, 16S rRNA gene cloning and *mcrA* (methyl coenzyme M reductase) gene cloning indicate the presence of methanogens (Table 3). In addition, the $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values for DR9IPC are similar to the values for microbial CH_4 identified in several other sites in the Witwatersrand Basin (Ward et al., 2004). In particular, fracture water from Masimong and Merriespruit, located 150 km southwest of Driefontein, have $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values for CH_4 of -60.7 and -207‰ (MM5); and -53.7 and -194‰ (MR1), respectively (Fig. 2). The microbial origin of CH_4 from Merriespruit is also supported by 16S rRNA gene and by enrichment cultures which indicate the presence of methanogens (Ward et al., 2004).

In contrast, 16S rRNA gene amplifications for the Kloof samples (KL739, KL441H2, KL443HWN1, KL443HWN) did not yield any evidence of methanogens (Ward et al., 2004; Kieft et al., 2005). While this




Fig. 2. Plot of $\delta^{13}\text{C}$ versus $\delta^2\text{H}$ values for CH_4 for sites from this study compared to the conventional fields for microbial and thermogenic CH_4 after Schoell (1988). Except for the two Mponeng samples (see text), samples fall along linear trends from the most ^{13}C -enriched end-members to more ^{13}C -depleted and ^2H -enriched end-members. Symbols are as in Fig. 1. DR938H3 and DR938H3' are samples taken from the same borehole 8 months apart. Unfortunately such temporal sampling was not possible for the other holes, due to the common practice of sealing them immediately after completion of drilling. Error bars are $\pm 0.5\text{‰}$ on $\delta^{13}\text{C}$ values and $\pm 5\text{‰}$ on $\delta^2\text{H}$ values and are smaller than the plotted symbols. Data from Masimong (MM) and Merriespruit (MR) are from Ward et al. (2004) for comparison. Data from Kidd Creek are from Sherwood Lollar et al. (2002).

may be due to the very low biomass in these samples and does not entirely rule out a small microbial CH_4 contribution, it is nonetheless consistent with the predominantly abiogenic origin proposed for CH_4 in the Kloof samples. No microbiological work was carried out at the other boreholes at Kloof, Copper Cliff South or Mponeng, but for Driefontein, 16S rRNA gene amplification and culture enrichments for methanogens were attempted for both borehole DR9IPC (results above) and for one of the proposed abiogenic end-members, borehole DR938H3. While enrichments were unsuccessful, PCR-amplified 16S rRNA gene indicated a bacterium (closely related to the sulfate-reducing *Desulfotomaculum* species) and three archaeal taxa within the family *Methanobacteriaceae*. The low biomass in this sample suggests that concentrations of CH_4 in the 100 μM range derived from microbial methanogenesis would be insufficient to isotopically overprint the $>17\text{ mM}$ abiogenic CH_4 signature however (Moser et al., in press).

The possibility of mixing between microbial hydrocarbons (characterized by ^{13}C -depleted and ^2H -enriched CH_4 values) and abiogenic end-members can be further evaluated using the traditional scheme shown in Fig. 3, adapted from Hunt (1996). In this plot, microbial hydrocarbons fall in the upper left corner based on their ^{13}C -depleted CH_4 values and CH_4/C_2^+ ratios >1000 (Hunt, 1996). We have modified the scheme to evaluate abiogenic-microbial mixing, using the abiogenic end-members for each site suggested by Fig. 1B. For the microbial

end-member, while microbial CH_4 has been identified at Kidd Creek and Copper Cliff South mines based on microbial cultures (Doig, 1994), no information on the isotopic composition of that CH_4 exists to date. Whereas a microbial end-member has been identified at Driefontein (DR9IPC) based on both isotopic evidence and gene cloning (Table 3), we do not have the same information for Kloof. The same dolomitic Transvaal Supergroup overlies Kloof mine however and as at Driefontein, fractures intersecting both the Ventersdorp and overlying Transvaal provide a conduit for mixing of fluids and gases between these two formations. For all samples then, as an initial test of the model, we used a $\delta^{13}\text{C}$ value of -55‰ for the microbial end-member based on the carbon isotope value of DR9IPC (Fig. 3). By this approach, data from Kloof are consistent with mixing of the most ^{13}C -enriched end-member (KL739) with up to 43% of a microbial hydrocarbon end-member with a $\delta^{13}\text{C}_{\text{CH}_4}$ value of -55‰ (Fig. 3). All estimates are insensitive to whether the microbial end-member has CH_4/C_2^+ ratios of 10^3 – 10^4 or even higher. For Driefontein, DR938H1 is consistent with mixing with up to 84% microbial CH_4 (based on mixing with DR548), while mixing between DR9IPC and DR938H3 defines a separate trend line, as in Fig. 2. Given that the gases occur in hydrogeologically isolated fractured rock it is not surprising that some sites exhibit a range of end-members and different possible mixing lines, as fracture controlled systems are rarely homogeneous. In fact, the mixing




Fig. 3. Plot of $\delta^{13}\text{C}$ values for CH_4 versus CH_4/C_2^+ ratios adapted from Hunt (1996). Data fall along mixing lines between the most ^{13}C -enriched CH_4 abiotic end-member at each site as suggested by Fig. 1B, and a ^{13}C -depleted CH_4 -rich microbial end-member, with a value of -55‰ (see text). Mixing lines are calculated based on the equations:

$$\delta^{13}\text{C} = x(\delta^{13}\text{C}_{\text{microbial}}) + (1 - x)(\delta^{13}\text{C}_{\text{abiotic}});$$

where x is the percent of the microbial end-member; and,

$$\text{CH}_4/\text{C}_2^+ = (x(I_{\text{microbial}}) + (1 - x)(I_{\text{abiotic}}))^{-1};$$

where $I = \text{C}_2^+/\text{CH}_4$, the inverse of CH_4/C_2^+ . The estimated percent microbial component for each sample (see text) is calculated by projecting its $\delta^{13}\text{C}$ value onto the mixing lines on Fig. 3. As noted in the text, the estimates of % microbial gas are relatively insensitive to CH_4/C_2^+ ratios but are dependent on the selected $\delta^{13}\text{C}$ values for the two end-members, and hence provide only a conservative estimate of the relative fraction of microbial gas. In particular, the mixing lines as depicted assume that the most ^{13}C -enriched end-member at each site is entirely free of a microbial contribution. If the most ^{13}C -enriched samples are already themselves the product of some mixing with microbial CH_4 with an even more ^{13}C -enriched end-member, then the estimates of the fraction of microbial contribution would be even larger. Symbols are as in Fig. 1. MM5 and MR1 (circles) are microbial CH_4 end-members previously identified for the Witwatersrand Basin based on CH_4 isotopic values, 16S rRNA gene and enrichment cultures (Ward et al., 2004). Unfortunately air contamination of sample DR9IPC means that C_2^+ values are diluted to below detection limit, so a CH_4/C_2^+ ratio cannot be calculated for this sample. Based on the similarity in isotopic composition of this sample to MR1 and MM5, it has been plotted (in brackets) close to those samples for comparison purposes. Error bars are $\pm 0.5\text{‰}$ on $\delta^{13}\text{C}$ values and $\pm 7\%$ on CH_4/C_2^+ ratios and are smaller than the plotted symbols.

lines could be recalculated for a microbial end-member with a $\delta^{13}\text{C}_{\text{CH}_4}$ value close to MM5 (-60‰) (Fig. 4) rather than close to DR9IPC (-55‰), and the data are still consistent with this simple two-component mixing between microbial and abiotic gas, although the estimates of the fraction of microbial mixing vary some-

what. The key point of Figs. 3 and 4 is not necessarily to constrain a specific microbial $\delta^{13}\text{C}$ value for each site, but to demonstrate that the isotopic and compositional variation in samples observed supports a model of mixing between the ^{13}C -enriched end-members in Fig. 1B, and microbial end-members with a range of $\delta^{13}\text{C}$ values typical of CH_4 produced by methanogens. If a similar approach is taken for the Canadian Shield sites, KC7792 can be explained by a mixture of the most ^{13}C -enriched end-member at Kidd Creek (KC8558), with up to 17.5% microbial CH_4 (Fig. 3). The CCS data fit somewhat less well. CCS4546 is the only sample with abiogenic characteristics at this site (Fig. 1B) and is the most ^{13}C -enriched and ^2H -depleted value at the site (Fig. 2). Hence, whereas the CH_4/C_2^+ ratio for CCS4546 in Fig. 3 is slightly elevated (8.6 versus approximately 6 for the other samples), a mixing line through the remainder of the data to a ^{13}C -enriched end-member with the $\delta^{13}\text{C}_{\text{CH}_4}$ value of CCS4546 and a CH_4/C_2^+ ratio of 6 indicates that a mixture of up to 23% microbial CH_4 can account for the rest of the data at that site.

The most ^{13}C -enriched end-members at each site are too C_2^+ -rich to be microbial in origin (Fig. 3), and likewise too ^2H -depleted in CH_4 to be consistent with a thermogenic origin (Fig. 2). Analogous to the Kidd Creek gases, the carbon and hydrogen isotopic patterns of these gases (Fig. 1B) suggest an abiogenic origin. The fact that the data from these sites in both Canada and South Africa fall on mixing lines in both Figs. 2 and 3 provides excellent support for interpretation of these data as two-component mixing between microbial




Fig. 4. Plot of $\delta^{13}\text{C}$ values for CH_4 versus CH_4/C_2^+ ratios adapted from Hunt (1996). In this plot, mixing lines are calculated as in Fig. 3 but for mixing between the most ^{13}C -enriched CH_4 end-member at each site, and a ^{13}C -depleted CH_4 rich microbial end-member with a value of -60‰ (see text).

and abiogenic end-members. Alternatively the presence of a thermogenic component at the sites in this study cannot be completely ruled out. Indeed, Ward et al. (2004) showed that mixing between microbial and thermogenic gases might account for gases in boreholes at Evander mine located in the quartzite, shale and minor volcanics of the Witwatersrand Supergroup sediments. Samples from South Africa in the present study however are predominantly from boreholes in a different geologic unit, the basaltic lavas of the Ventersdorp Supergroup. Furthermore, all the samples at Evander fall close to the conventional microbial and thermogenic isotope fields in Fig. 2 (Ward et al., 2004), unlike the ^{13}C -enriched and ^2H -depleted end-members at the sites in the present study. Hence we feel the best explanation for the data in the present study is indeed two-component mixing between ^{13}C -enriched abiogenic and ^{13}C -depleted microbial end-members. The fact that data from this study fit a simple two-component mixing model between these microbial and abiogenic end-members suggests that if a thermogenic component is present, it is small and does not control the isotopic and compositional variations in a major way.

4.3. H_2 -based autotrophy linked to methanogenesis

This approach to resolving mixing between abiogenic and microbial CH_4 in the crust has important implications for evolving research into the Earth's deep biosphere, specifically the identification of potential substrates supporting microbial communities in the deep subsurface. H_2 -based subsurface microbial communities associated with autotrophic methanogenesis have been proposed in the Columbia River basalt aquifer (Anderson et al., 1998; Stevens and McKinley, 1995), Lidy Hot springs (Chapelle et al., 2002); and in carbonate-rich serpentinized peridotites at an off-axis hydrothermal vent field near the mid-Atlantic ridge (Kelley et al., 2001; 2005). Interest in such systems is high since they may be modern terrestrial analogues for subsurface microbial ecosystems under reducing conditions on the early Earth (Shock and Schulte, 1998) or on other planets or Jovian satellites (Chapelle et al., 2002; McCollom, 1999). Several of the boreholes sampled in this study have high concentrations of H_2 gas as well as the CH_4 and higher hydrocarbons already discussed. H_2 concentrations in samples from Mponeng, Driefontein and Kloof are as high as 11.5% (Table 1). H_2 levels in samples from CCS, Kidd Creek and other sites in the Precambrian Shield rocks of Canada have been reported in concentrations as high as 1–58% (Table 1; (Sherwood Lollar et al., 1993b)). Over several

years, re-sampling of boreholes in Canada has shown that the high H_2 levels are not simply an artefact of drilling (Bjerg et al., 1997) but a persistent natural phenomenon in this geologic environment (Sherwood et al., 1988; Sherwood Lollar et al., 1993b, 2002). Like the hydrocarbon gases, the high quantities of H_2 are likely the product of abiogenic water–rock interactions in these geologically isolated, high rock to water ratio environments. Specific H_2 -generating reactions may vary from one geological environment to another. At sites such as Kidd Creek where significant sections of ultramafic rock occur, H_2 may be the product of serpentinization over geologically long time scales analogous to systems described elsewhere (Abrajano et al., 1988; Charlou and Donval, 1993; Kelley et al., 2001). In the absence of significant ultramafics, such as at the Witwatersrand Basin sites, radiolytic decomposition of water in the uranium-rich Witwatersrand Formation has been shown to produce H_2 at a sufficient rate to account for the elevated H_2 concentrations observed in these isolated fracture fluids (Lin et al., 2005). H_2 as an energy source for autotrophic microbial ecosystems on both the Earth (Anderson et al., 1998; Stevens and McKinley, 1995) and Mars (Boston et al., 1992) has been a topic of considerable debate in the literature. Nonetheless, the number of studies that have been able to convincingly demonstrate coupled oxidation of H_2 and reduction of CO_2 to produce CH_4 via the equation:



as the basis for a microbial food chain in the deep terrestrial subsurface has been limited (Chapelle et al., 2002).

The approach for identifying abiogenic and microbial CH_4 end-members presented in this paper, and the large database on deep subsurface gases developed by our research in Canada and South Africa provides an opportunity to test this proposed reaction on a large scale. Chemolithoautotrophs growing on H_2 can potentially reduce a variety of different oxidants including any available oxygen, sulfur and nitrogen oxidants, metals and metalloids, as well as CO_2 . If H_2 autotrophy linked to CO_2 reduction and methanogenesis is an important process, there may be a correlation between H_2 utilization and production of an isotopically ^{13}C -depleted microbial CH_4 . Fig. 5 is a compilation of all samples for Canada and South Africa measured by our group to date ($n=65$), incorporating data from almost two dozen sites. A relationship exists between H_2 concentrations and the isotopically distinct microbial and abiogenic hydrocarbon gas end-members identified in this study. With few excep-




Fig. 5. Histogram of H_2 concentrations versus $\delta^{13}\text{C}_{\text{CH}_4}$ values for samples ($n=65$) from this study as well as for previously reported data from Precambrian Shield sites in South Africa (Ward et al., 2004) and Canada (Montgomery, 1994; Sherwood Lollar et al., 1993a,b, 2002). With few exceptions, high H_2 concentrations (>1%) ($n=31$) are associated with the most ^{13}C -enriched CH_4 end-members ($\delta^{13}\text{C}_{\text{CH}_4}$ values between -45% and -25%). In contrast, samples with ^{13}C -depleted CH_4 values (indicative of a significant methanogenic component) have H_2 values between 0.1–1% ($n=7$), to below detection limit (<0.01%) ($n=27$).

tions, high H_2 concentrations (>1%) are associated with only the most ^{13}C -enriched CH_4 samples ($\delta^{13}\text{C}$ values between -45 and -25%). In contrast, samples with ^{13}C -depleted CH_4 values (indicative of a significant methanogenic component) correlate with H_2 values in the range of 0.1% to 1%, to below detection limit (<0.01%). The observed pattern of H_2 concentrations and CH_4 isotopic signatures may be the geochemical fingerprint of H_2 autotrophy linked to methanogenesis via Eq. (1). This evidence supports the interpretation of the hydrocarbon isotopic data as controlled by two-component mixing between ^{13}C -enriched CH_4 end-members and ^{13}C -depleted microbial end-members and may further suggest that methanogenesis is supported by H_2 utilization. A scenario could be envisaged where H_2 – CH_4 rich abiogenic gases are mobilized when fractures open due to tectonic activity or due to more recent disturbances due to mining activity and support a redox gradient-driven ecosystem in the Precambrian Shield deep subsurface where rapid utilization of H_2 by microbial communities quickly depletes H_2 and produces a ^{13}C -depleted CH_4 signature characteristic of methanogenesis that mixes with and partially overprints the pre-existing ^{13}C -enriched abiogenic CH_4 signature.

5. Conclusions

This study suggests that the abiogenic hydrocarbon gases first identified at Kidd Creek in fact exist at a variety of Precambrian Shield sites worldwide. Using the mixing models outlined here, the relative contribution of abiogenic versus microbial end-members can be estimated. Verification of an abiogenic component first requires identification of the end-member based on the $\delta^{13}\text{C}$ and $\delta^2\text{H}$ model illustrated in Fig. 1, followed by evaluation of the extent of mixing with other more conventional hydrocarbon sources via mixing schemes such as in Figs. 2 and 3. This combined approach has important implications for evaluation of potential abiogenic hydrocarbon reservoirs in a wide range of geologic settings, including the longstanding controversy concerning the possible contribution of abiogenic gases to economic petroleum hydrocarbon reservoirs (Gold, 1979; Kenney et al., 2002). The association of high concentrations of H_2 with ^{13}C -enriched CH_4 end-members, and the H_2 depletion in ^{13}C -depleted methanogenic end-members further suggests the possibility that abiogenic gases may support H_2 autotrophy linked to methanogenesis in the deep subsurface.

Acknowledgements

This study was supported in part by grants from the Natural Sciences and Engineering Research Council of Canada and Discovery Grants Program, and Canadian Space Agency to BSL and by the National Science Foundation Life in Extreme Environments Program (EAR-9714214) grant to TCO and NASA Astrobiology Institute grant to the IPTAI team. We thank H. Li for compositional and isotopic analyses. Special thanks are due to the geologists and staff of the following mines for providing geological information and invaluable assistance with underground field work: Kidd Creek Mine, Timmins, Ontario (D. Duff, R. Cook, P. Olsen); Copper Cliff South Mine, Sudbury, Ontario (T. Little); and in South Africa, Driefontein (D. Nel, B. Kotze, M. deKoker), Mponeng (D. Kirshaw, H. Cole, G. Gilchrest) and Kloof (A. Van Heerden, D. Steinkamp, T. Hewitt, G. Buxton) Mines. Special thanks are owed to Rob Wilson of SRK-Turgis for logistical support. [SG]

References

- Abrajano, T.A., Sturchio, N.C., Bohlke, J.K., Lyon, G.L., Poreda, R.J., Stevens, C.M., 1988. Methane–hydrogen gas seeps, Zambales Ophiolite, Philippines: deep or shallow origin? *Chemical Geology* 71, 211–222.

- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J., 1997. Gapped BLAST and PSA-BLAST: a new generation of protein database search program. *Nucleic Acids Research* 25, 3389–3402.
- Anderson, R.B., 1984. The Fischer–Tropsch Synthesis. Academic Press.
- Anderson, R.T., Chapelle, F.H., Lovley, D.R., 1998. Evidence against hydrogen-based microbial ecosystems in basalt aquifers. *Science* 281 (5379), 976–977.
- Baker, B.J., Moser, D.P., MacGregor, B.J., Fishbain, S., Wagner, M., Fry, N.K., Jackson, B., Speolstra, N., Loos, S., Takai, K., Sherwood Lollar, B., Fredrickson, J.K., Balkwill, D., Onstott, T.C., Wimpee, C.F., Stahl, D.A., 2003. Related assemblages of sulphate-reducing bacteria associated with ultradeep gold mines of South Africa and deep basalt aquifers of Washington State. *Environmental Microbiology* 5 (4), 267–277.
- Berndt, M.E., Allen, D.E., Seyfried, W.E.J., 1996. Reduction of CO₂ during serpentinization of olivine at 300 °C and 500 bar. *Geology* 24 (4), 351–354.
- Bjerg, P.L., Jakobsen, R., Bay, H., Rasmussen, M., Albrechtsen, H.-J., Christensen, T.H., 1997. Effects of sampling well construction on H₂ measurements made for characterization of redox conditions in a contaminated aquifer. *Environmental Science and Technology* 31, 3029–3031.
- Boston, P.M., Ivanov, M.V., McKay, C.P., 1992. On the possibility of chemosynthetic ecosystems in subsurface habitats on Mars. *Icarus* 95, 300–308.
- Chapelle, F.H., O'Neill, K., Bradley, P.M., Methe, B.A., Ciufo, S.A., Knobel, L.L., Lovley, D.R., 2002. A hydrogen-based subsurface microbial community dominated by methanogens. *Nature* 415, 312–315.
- Charlou, J.L., Donval, J.P., 1993. Hydrothermal methane venting between 12N and 26N along the Mid-Atlantic ridge. *Journal of Geophysical Research* 98, 9625–9642.
- Cochrane, L.B., 1984. Ore deposits of the Copper Cliff offset. In: Kendrick, G. (Ed.), *The Geology and Ore deposits of the Sudbury Structure*. Ontario Geological Survey, pp. 347–360.
- Cook, A.P., 1998. The occurrence, emission and ignition of combustible strata gases in Witwatersrand gold mines and Bushveld platinum mines and means of ameliorating related ignition and explosion hazards. Safety in Mines Research Advisory Committee, Interim Project Report, vol. 504. Itasca Africa (Pty) Ltd. 90 pgs.
- Coward, M.P., Spencer, R.M., Spencer, C.E., 1995. Development of the Witwatersrand Basin, South Africa. *Early Precambrian Processes* 95, 243–269.
- Des Marais, D.J., Donchin, J.H., Nehring, N.L., Truesdell, A.H., 1981. Molecular carbon isotopic evidence for the origin of geothermal hydrocarbons. *Nature* 292, 826–828.
- Doig, F., 1994. Bacterial methanogenesis in Canadian Shield groundwaters. M.Sc. Thesis, University of Toronto, Toronto. 99 pp.
- Formisano, V., Atreva, S., Encrenaz, T., Ignatiev, N., Giuranna, M., 2004. Detection of methane in the atmosphere of Mars. *Science* 306, 1758–1761.
- Foustoukos, D.I., Seyfried, W.E.J., 2004. Hydrocarbons in hydrothermal vent fluids: the role of chrome-bearing catalysts. *Science* 304, 1002.
- Giardini, A.A., Salotti, C.A., 1969. Kinetics and relations in the calcite–hydrogen reaction and relations in the dolomite–hydrogen and siderite–hydrogen systems. *American Mineralogist* 54, 1151–1172.
- Gold, T., 1979. Terrestrial sources of carbon and earthquake outgassing. *Journal of Petroleum Geology* 1, 1–19.
- Hales, B.A., Edwards, C., Ritchie, D.A., Hall, G., Pickup, R.W., Saunders, J.R., 1996. Isolation and identification of methanogen-specific DNA from blanket bog peat by PCR amplification and sequence analysis. *Applied and Environmental Microbiology* 62, 668–675.
- Holloway, J.R., 1984. Graphite–CH₄–H₂O–CO₂ equilibria at low grade metamorphic conditions. *Geology* 12, 455–458.
- Horita, J., Berndt, M.E., 1999. Abiogenic methane formation and isotopic fractionation under hydrothermal conditions. *Science* 285, 1055–1057.
- Hu, G., Ouyang, Z., Wang, X., Wen, Q., 1998. Carbon isotope fractionation in the process of Fischer–Tropsch reaction in primitive solar nebula. *Science China* 41, 202–207.
- Hunt, J.M., 1996. *Petroleum Geochemistry and Geology*. W.H. Freeman and Company, New York. 743 pp.
- Jenden, P.D., Kaplan, I.R., 1989. Origin of natural gas in the Sacramento Basin, California. *American Association of Petroleum Geologists Bulletin* 73, 431–453.
- Jenden, P.D., Kaplan, I.R., Poreda, R.J., Craig, H., 1988. Origin of nitrogen-rich natural gases in the California Great Valley: evidence from helium, carbon and nitrogen isotope ratios. *Geochimica et Cosmochimica Acta* 52, 851–861.
- Jenden, P.D., Hilton, D.R., Kaplan, I.R., Craig, H., 1993. Abiogenic hydrocarbons and mantle helium in oil and gas fields. In: Howell, D.G. (Ed.), *The Future of Energy Gases*, U.S. Geological Survey Professional Paper, vol. 1570, pp. 31–56.
- Kelley, D.S., Karson, J.A., Blackman, D.K., Fruh-Green, G.L., Butterfield, D.A., Lilley, M.D., Olson, E.J., Schrenk, M.O., Roe, K.K., Lebon, G.T., Rivizzigno, P., 2001. An off-axis hydrothermal vent field near the Mid-Atlantic Ridge at 30°N. *Nature* 412, 145–149.
- Kelley, D.S., Karson, J.A., Fruh-Green, G.L., Yoerger, D.R., Shank, T.M., Butterfield, D.A., Hayes, J.M., Schrenk, M.O., Olson, E.J., Proskurowski, G., Jakuba, M., Bradley, A., Larson, B., Ludwig, K., Glickson, D., Buckman, K., Bradley, A.S., Brazelton, W.J., Roe, K., Elend, M.J., Delacour, A., Bernasconi, S.M., Lilley, M.D., Baross, J.A., Summons, R.T., Sylva, S.P., 2005. A serpentine-hosted ecosystem: the Lost City Hydrothermal Field. *Science* 307, 1428–1434.
- Kenney, J.F., Kutcherov, V.A., Bendeliani, N.A., Alekseev, V.A., 2002. The evolution of multicomponent systems at high pressures: the thermodynamic stability of the hydrogen-carbon system: the genesis of hydrocarbons and the origin of petroleum. *PNAS* 99 (17), 10976–10981.
- Kerr, R.A., 2004. Heavy breathing on Mars? *Science* 306, 29.
- Kieft, T.L., Fredrickson, J.K., Onstott, T.C., Gorby, Y.A., Kostandarithes, H.M., Bailey, T.J., Kennedy, D.W., Li, S.W., Plymale, A.E., Spadoni, C.M., Gray, M.S., 1999. Dissimilatory reduction of Fe(III) and other electron acceptors by a *Thermus* isolate. *Applied and Environmental Microbiology* 65 (3), 1214–1221.
- Kieft, T.L., McCuddy, S.M., Onstott, T.C., Davidson, M., Lin, L.-H., Mislowack, B., Pratt, L., Boice, E., Sherwood Lollar, B., Lippmann-Pipke, J., Pfiffner, S.M., Phelps, T.J., Gihring, B., Moser, D., van Heerden, A., 2005. Geochemically generated, energy-rich substrates and indigenous microorganisms in deep, ancient groundwater. *Geomicrobiology Journal* 22, 325–335.
- Lahermo, P.W., Lampen, P.H., 1987. Brackish and saline groundwaters in Finland. In: Fritz, P., Frapé, S.K. (Eds.), *Saline Water and Gases in Crystalline Rock*, Geological Association of Canada Special Paper, vol. 33, pp. 103–110.
- Lin, L.-H., Slater, G.F., Sherwood Lollar, B., Lacrampe-Couloume, G., Onstott, T.C., 2005. The yield and isotopic composition of

- radiolytic H₂, a potential energy source for the deep subsurface biosphere. *Geochimica et Cosmochimica Acta* 69, 893–903.
- Lippmann, J., Stute, M., Torgersen, T., Moser, D.P., Hall, J., Lin, L., Borcsik, M., Bellamy, R.E.S., Onstott, T.C., 2003. Dating ultra-deep mine waters with noble gases and ³⁶Cl, Witwatersrand Basin, South Africa. *Geochimica et Cosmochimica Acta* 67, 4597–4619.
- Lyon, G.L., Hulston, J.R., 1984. Carbon and hydrogen isotopic compositions of New Zealand geothermal gases. *Geochimica et Cosmochimica Acta* 48, 1161–1171.
- McCollom, T.M., 1999. Methanogenesis as a potential source of chemical energy for primary biomass production by autotrophic organisms in hydrothermal systems on Europa. *Journal of Geophysical Research* 104 (E12), 30729–30742.
- McCollom, T.M., Seewald, J.S., 2001. A reassessment of the potential for reduction of dissolved CO₂ to hydrocarbons during serpentinization of olivine. *Geochimica et Cosmochimica Acta* 65 (21), 3769–3778.
- Montgomery, J., 1994. An isotopic study of CH₄ and associated N₂ and H₂ gases in Canadian Shield mining environments. MSc Thesis, University of Toronto, Toronto, Canada, 72 pp.
- Moser, D.P., Gihring, T.M., Brockman, F.J., Fredrickson, J.K., Balkwill, D.L., Dollhopf, M.E., Sherwood Lollar, B., Pratt, L.M., Boice, E., Southam, G., Wander, G., Baker, B.J., Pfiffner, S.M., Lin, L.-H., Onstott, T.C., 2005. Deep continental fracture system dominated by *Methanobacterium* spp. and *Desulfotomaculum* spp. *Applied and Environmental Microbiology*. In press.
- Moyer, C.L., Tiedje, J.M., Dobbs, F.C., Karl, D.M., 1998. Diversity of deep-sea hydrothermal vent Archaea from Loihi Seamount, Hawaii. *Deep-Sea Research II* 45, 303–317.
- Nurmi, P.A., Kukkonen, I.T., 1986. Geochemistry of water and gas from deep drill holes: a new sampling technique. *Canadian Journal of Earth Sciences* 23, 1450–1454.
- Onstott, T.C., Moser, D.P., Pfiffner, S.M., Fredrickson, J.K., Brockman, F.J., Phelps, T.J., White, D.C., Peacock, A., Balkwill, D., Hoover, R., Krumholz, L.R., Borcsik, M., Kieft, T.L., Wilson, R., 2003. Indigenous and contaminant microbes in ultradeep mines. *Environmental Microbiology* 5, 1168–1191.
- Oremland, R.S., Des Marais, D.J., 1983. Distribution, abundance and carbon isotopic composition of gaseous hydrocarbons in Big Soda Lake, Nevada: an alkaline, meromictic lake. *Geochimica et Cosmochimica Acta* 47, 2107–2114.
- Poreda, R.J., Jenden, P.D., Kaplan, I.R., Craig, H., 1986. Mantle helium in Sacramento basin natural gas wells. *Geochimica et Cosmochimica Acta* 50, 2847–2853.
- Poreda, R.J., Jeffrey, A.W.A., Kaplan, I.R., Craig, H., 1988. Magmatic helium in subduction-zone natural gases. *Chemical Geology* 71, 199–210.
- Reysenbach, A.L., Longnecker, K., Kirshtein, J., 2000. Novel bacterial and archaeal lineages from an in situ growth chamber deployed at a Mid-Atlantic Ridge hydrothermal vent. *Applied and Environmental Microbiology* 66, 3798–3806.
- Rigby, D., Smith, J.W., 1981. An isotopic study of gases and hydrocarbons in the Cooper Basin. *Australian Petroleum Exploration Association Journal* 21, 222–229.
- Schoell, M., 1988. Multiple origins of methane in the Earth. *Chemical Geology* 71, 1–10.
- Sherwood, B., Fritz, P., Frappe, S.K., Macko, S.A., Weise, S.M., Welhan, J.A., 1988. Methane occurrences in the Canadian Shield. *Chemical Geology* 71, 223–236.
- Sherwood Lollar, B., Frappe, S.K., Fritz, P., Macko, S.A., Welhan, J.A., Blomqvist, R., Lahermo, P.W., 1993a. Evidence for bacterially generated hydrocarbon gas in Canadian Shield and Fennoscandian Shield rocks. *Geochimica et Cosmochimica Acta* 57, 5073–5085.
- Sherwood Lollar, B., Frappe, S.K., Weise, S.M., Fritz, P., Macko, S.A., Welhan, J.A., 1993b. Abiogenic methanogenesis in crystalline rocks. *Geochimica et Cosmochimica Acta* 57, 5087–5097.
- Sherwood Lollar, B., Frappe, S.K., Weise, S.M., 1994. New sampling devices for environmental characterization of groundwater and dissolved gas chemistry (CH₄, N₂, He). *Environmental Science and Technology* 28 (13), 2423–2427.
- Sherwood Lollar, B., Westgate, T.D., Ward, J.A., Slater, G.F., Lacrampe-Couloume, G., 2002. Abiogenic formation of gaseous alkanes in the Earth's crust as a minor source of global hydrocarbon reservoirs. *Nature* 416, 522–524.
- Shock, E.L., 1995. An open or shut case? *Nature* 378, 338–339.
- Shock, E.L., Schulte, M.D., 1998. Organic synthesis during fluid mixing in hydrothermal systems. *Journal of Geophysical Research* 103 (E12), 28513–28527.
- Stevens, T.O., McKinley, J.P., 1995. Lithoautotrophic microbial ecosystems in deep basalt aquifers. *Science* 270, 450–454.
- Takai, K., Moser, D.P., DeFlaun, M., Onstott, T.C., Fredrickson, J.K., 2001. Archaeal diversity in waters from deep South African gold mines. *Applied and Environmental Microbiology* 67 (12), 5750–5760.
- Vanko, D.A., Stakes, D.S., 1991. Fluids in oceanic layer 3: evidence from veined rocks, hole 735B, Southwest Indian Ridge. *Proceedings of the Oceanic Drilling Program. Scientific Results* 118, 181–215.
- Ward, J.A., Slater, G.F., Moser, D.P., Lin, L.-H., Lacrampe-Couloume, G., Bonin, A.S., Davidson, M., Hall, J.A., Mislouack, B.J., Bellamy, R.E.S., Onstott, T.C., Sherwood Lollar, B., 2004. Microbial hydrocarbon gases in the Witwatersrand Basin, South Africa: implications for the deep biosphere. *Geochimica et Cosmochimica Acta* 68 (13), 3239–3250.
- Yuen, G.U., Pecore, J.A., Kerridge, J.F., Pinnavaia, T.J., Rightor, E.G., Flores, J., Wedeking, K., Mariner, R., Des Marais, D.J., Chang, S., 1990. Carbon isotope fractionation in Fischer-Tropsch type reactions. *Lunar Planetary Science Conference XXI*, pp. 1367–1368.